

Ecological Study of Dietary and Smoking Links to Lymphoma

William B. Grant, PhD

Abstract

The ecological approach is used to investigate dietary and smoking links to lymphoma. International mortality rate data for 1986 and 1994 by gender and age group are compared with national dietary supply values of various food components for up to 10 years prior to the mortality data as well as per capita cigarette consumption rates 5 and 15 years earlier. The non-fat portion of milk, 3-9 years prior to the 1986 mortality data and 4 years prior to the 1994 data, was found to have the highest association with lymphoma, with r as high as 0.89. The results imply that 70 percent of lymphoma mortality may be related to this dietary component. Cigarette smoking in 1980 was found to have a weaker association with 1994 lymphoma mortality rates, being most important for younger men and statistically insignificant for younger women. The non-fat milk result is consistent with both case-control studies and a Norwegian prospective study, and with the often-observed finding that abnormal calcium metabolism, hypercalciuria, and dysregulated calcitriol production are common in normocalcemic patients with non-Hodgkin's lymphoma (NHL). It is hypothesized that excess dietary calcium from milk is a significant risk factor for lymphoma.

(*Altern Med Rev* 2000;5(6):563-562)

Introduction

A number of environmental, dietary, and lifestyle factors are involved in the etiology of lymphoma. Various chemicals, including herbicides^{1,2} and pesticides,³ have been found to be associated with increased risk. Occupations with significant chemical exposure, such as agricultural work,⁴⁻⁶ as well as smoking^{7,8} have been determined to be risk factors for lymphoma. There are a number of foods which have been associated with an increased risk for lymphoma. Several studies, using either prospective cohort⁹ or case-control approaches,¹⁰⁻¹² have found milk to be a risk factor. However, one study found animal fat to be the highest risk factor (risk ratio (RR) = 2.00) in older women, with milk and dairy product consumption not associated with NHL.¹³ A study in Uruguay reported that salted meat was a high risk factor (odds ratio (OR) = 5.5).⁸ A U.S. study reported that red meat (multivariate relative risk (MRR) = 2.2) and trans unsaturated fat (MRR = 2.4) were risk factors.¹⁴ Finally, a lectin found in raw kidney bean, phytohaemagglutinin (PHA), was found to *reduce* the rate of an NHL tumor in mice.¹⁵

William B. Grant, PhD – Lidar Applications Group, Atmospheric Sciences Research, NASA Langley Research Center. In his spare time he has taken a special interest in examination of the epidemiological relationship between diet and disease. Correspondence address: 12 Sir Francis Wyatt Place, Newport News, VA 23606-3660. Email: wbgrant@norfolk.infi.net

Table 1: Data for Lymphoma Mortality for Males and Females Aged 65-74 Years for 1986 and 1994 along with Non-fat Milk Supply for 1977 and 1988.

| Country | 1986/1994 Mortality Rate | | 1977/1988 Non-fat Milk |
|----------------|--------------------------|-----------|------------------------|
| | males | females | kcal/day/person |
| | cases/100,000/year | | |
| Argentina | 25.9/29.2 | 19.0/19.7 | 105/116 |
| Australia | 56.4/56.6 | 38.2/44.0 | 186/206 |
| Austria | 33.7/43.9 | 27.1/30.3 | 157/166 |
| Belgium | 34.5/42.3 | 26.3/26.7 | 153/146 |
| Canada | 56.8/56.4 | 37.8/39.0 | 155/141 |
| Chile | 11.2/39.6 | 13.3/22.6 | 85/74 |
| Columbia | --/25.5 | --/18.1 | --/76 |
| Costa Rica | 30.5/24.7 | 32.5/22.5 | 126/120 |
| Cuba | 37.7/35.3 | 30.0/25.1 | 154/146 |
| Czechoslovakia | 40.1/-- | 25.0/-- | 147/-- |
| Denmark | 52.4/56.2 | 37.9/28.4 | 159/162 |
| Equador | 10.6/-- | 7.1/-- | 24/-- |
| Finland | 51.6/60.7 | 34.1/33.5 | 238/202 |
| France | 39.7/43.7 | 23.3/28.2 | 150/176 |
| Germany | 36.2/42.3 | 22.5/27.7 | 134/151 |
| Greece | 23.6/24.8 | 14.3/14.4 | 129/130 |
| Hong Kong | 21.4/29.0 | 16.8/14.3 | 50/59 |
| Hungary | 34.7/42.4 | 25.7/25.4 | 99/114 |
| Ireland | 39.3/49.9 | 38.1/39.9 | 182/253 |
| Italy | 36.7/44.7 | 24.9/31.5 | 134/113 |

population was less than two million, if the mortality rates for those aged 75+ years were lower than for those aged 65-74 years, or if the countries were part of the former Soviet Union because these countries do not have medical care systems on par with those in the West. Israel was also omitted because of the large influx of immigrants in recent history. See Figure 1 for 1986 lymphoma mortality rates for males aged 75+ years compared to 1980 non-fat milk consumption. Figure 2 depicts 1994 lymphoma mortality rates for females aged 65-74 years versus 1993 non-fat milk supply.


National dietary supply data for this study were obtained from the Food and Agriculture Organization (FAO) *Food Balance Sheets*.²⁰ Such data are often used in international ecologic studies. It has been shown that in the United States 75 percent of the consumer food supply is actually consumed by people.²² It is assumed that similar factors apply to other countries. FAO dietary fat data appear to be consistent with studies from various European countries.²³ To further examine the usefulness of the FAO dietary supply values for this study,

It should also be noted that neither childhood leukemia nor lymphoma have strong dietary links but seem to be linked to infections.¹⁶⁻¹⁸

Methods

Lymphoma mortality data used in this study were obtained for 38 countries for 1986 and 33 countries for 1994 (+1/-2 years) from the World Health Organization's *World Health Statistics Annual*.¹⁹ The countries used in the analysis along with the 1986 and 1994 mortality rates for those aged 65-74 years are given in Table 1. Countries were excluded if the

Figure 1: 1986 Lymphoma Mortality Rates, Males (aged 75+ years), vs. 1980 Non-fat Milk Supply.




those values are compared with recent surveys of food intakes in ten Western European countries.²⁴ Aside from alcohol, the energy values in the studies compared to the FAO data for carbohydrates, total energy, fat, and protein lie in the range from 0.62 to 0.88. The value of 0.62 for fat when the Mediterranean countries are included could be due to widespread use of olive oil in cooking not reflected in intake. The study values are also affected by underreporting²⁵ and, in the case of alcohol, whether the study was conducted during the week or weekend.²⁶ Nonetheless, the standard deviation (SD) of the values in terms of energy, omitting alcohol, lie in the 0.12-0.20 range. The SDs for the primary categories as fractions of total energy lie in the range 0.04-0.12. The statistical results for dietary components and disease values are similar whether absolute or energy-normalized dietary values

are used. These SD values are much lower than the variations of the FAO data between countries. Thus, it appears appropriate to assume the FAO data can be used in such ecological studies.

A number of dietary components were used in the analysis, including animal fat, cereals, dairy fat, fruit, milk, total fat, total energy, and vegetable products. The years used in the analysis for dietary components extended from 1977-1983 for the 1986 mortality data, and 1988-1993 for the 1994 mortality data, since it cannot be determined *a priori* which years are most important in the etiology. Linear and multiple linear regression analyses were performed on the various data sets. The F-value was used to rank the results and evaluate the multiple linear regression results. Smoking data were obtained²⁷ and reported as number of cigarettes smoked per

Figure 2: 1994 Lymphoma Mortality Rates, Females (aged 65-74 years), vs. 1993 Non-fat Milk Supply.



person per year for 1980-1982 and 1990-1992.

Results

The highest statistical association between dietary factors and lymphoma was found for the non-fat portion of milk (Tables 2 and 3). The non-fat portion of milk is calculated from total energy derived from milk minus four times the milk protein weight (grams) and nine times the milk fat weight (grams). The statistical results passed all tests applied, including normality, constant variance, and test with alpha.

Cigarette smoking per capita for 1980 also showed a positive association with lymphoma mortality rates, being highest for males aged 55-64 years, and lowest for females aged 55-64 years (Table 4). However, the inclusion

of cigarette smoking in the multiple linear regression reduced, rather than increased, the F (t^2 , student-t test) value, indicating that smoking as measured in that manner had a weak association with lymphoma mortality. While other dietary components had positive or inverse associations with lymphoma mortality, none increased the F value in a multiple linear regression, and none did as well as cigarette smoking, although cereals came close. Therefore, the ecological approach cannot determine whether they are risk or risk-reduction factors or merely confounding factors, so these results are not reported here.

Table 2: Statistical Results of Lymphoma Using 1986 Mortality Data and the Non-fat Portion of Milk for Various Years for 38 Countries.

| Gender, Age | Year of dietary data | | |
|---------------|----------------------|------------|------------|
| | 1977 | 1980 | 1983 |
| Years | r, F | r, F | r, F |
| MALE, 55-64 | 0.820, 74 | 0.796, 62 | 0.815, 71 |
| FEMALE, 55-64 | 0.786, 56 | 0.763, 49 | 0.767, 50 |
| MALE, 65-74 | 0.817, 72 | 0.807, 67 | 0.831, 81 |
| FEMALE, 65-74 | 0.840, 86 | 0.817, 72 | 0.816, 77 |
| MALE, 75+ | 0.875, 117 | 0.868, 110 | 0.867, 109 |
| FEMALE, 75+ | 0.885, 130 | 0.880, 124 | 0.881, 125 |

Table 3: Statistical Results for Lymphoma Using 1994 Mortality Data and the Non-fat Portion of Milk from Various Years for 33 Countries.

| Gender, Age | Year of dietary data | | |
|---------------|----------------------|-----------|-----------|
| | 1988 | 1990 | 1993 |
| Years | r, F | r, F | r, F |
| MALE, 55-64 | 0.688, 28 | 0.691, 28 | 0.654, 23 |
| MALE, 65-74 | 0.811, 60 | 0.827, 67 | 0.782, 49 |
| MALE, 75+ | 0.853, 83 | 0.863, 91 | 0.833, 70 |
| FEMALE, 55-64 | 0.752, 40 | 0.736, 37 | 0.637, 21 |
| FEMALE, 65-74 | 0.870, 97 | 0.863, 90 | 0.870, 97 |
| FEMALE, 75+ | 0.830, 69 | 0.847, 79 | 0.816, 62 |

Discussion

The statistical results for adult lymphoma reported here are consistent with several prior studies in which milk was found to be associated with mortality from the disease. In a prospective cohort study in Norway,⁹ cancer of the lymphatic organs was the cancer found to have the highest association with milk consumption (OR for >2 glasses/day vs <1 glass/day was 3.36, 95% CI = 1.4-8.2). In a case-control study of dietary factors and NHL,¹⁰ milk had the highest statistical association. However, one study found an insignificantly elevated risk of NHL among men with high milk consumption.¹¹ A case-control study¹² reported an OR for milk for the highest tertile of 1.8 for NHL. They also reported an OR for whole-grain foods of 0.4. It is not clear why other reports have failed to find a connection between milk consumption and NHL in women.¹³ However, their analysis did not consider the non-fat portion of milk separately, nor the amount of cereals in the diet. They did find reduced RRs for carbohydrates and dietary fiber.

The non-fat milk supply values for 1977²⁰ and 1988²¹ are given in Table 1. For the 17 countries for which non-fat milk increased, lymphoma mortality rates increased for males and/or females aged 65-74 years in 14 countries. There are six countries for which Table 1 shows a decrease in non-fat milk supply while another source²¹ showed either no change or a slight increase. Excluding these countries leaves eight for which the non-fat milk supply clearly decreased. For these countries, lymphoma mortality rates for males aged 65-74 years decreased in four of the countries, while they decreased for females aged 65-74 years in three countries. Some of the

Table 4: Statistical Results for Lymphoma Mortality for 32 Countries in 1994 Including Non-fat Milk for 1990 and Cigarettes per capita for 1980 (risk factors) and Cereals (risk reduction factor).

| Gender | Age | r, F | p(non-fat milk) | p(cigarettes) | p(cereals) |
|--------|-------|-----------|-----------------|---------------|------------|
| Male | 55-64 | 0.774, 22 | <0.001 | 0.005 | |
| | 65-74 | 0.849, 38 | <0.001 | 0.060 | |
| | 75+ | 0.889, 55 | <0.001 | 0.060 | |
| Female | 55-64 | 0.738, 17 | <0.001 | 0.303 | |
| | 65-74 | 0.877, 48 | <0.001 | 0.068 | |
| | 75+ | 0.869, 45 | <0.001 | 0.049 | |
| Male | 55-64 | 0.774, 14 | 0.004 | 0.006 | 0.900 |
| | 65-74 | 0.856, 26 | <0.001 | 0.072 | 0.289 |
| | 75+ | 0.892, 36 | <0.001 | 0.071 | 0.470 |
| Female | 55-64 | 0.742, 11 | 0.005 | 0.333 | 0.525 |
| | 65-74 | 0.879, 32 | <0.001 | 0.079 | 0.560 |
| | 75+ | 0.886, 34 | <0.001 | 0.057 | 0.058 |

countries, such as Chile, may be seeing an increase in lymphoma mortality rates due to increased use of agricultural chemicals.

The ratios of lymphoid mortality rates for males to females for three age groups are listed in Table 5. There is no statistically significant gender difference. This suggests that gender-specific hormones are not involved, since for other diseases such as acute myocardial infarction, female hormones seem to play a role, resulting in rapid ratio changes above 55 years of age.²⁸

Since adult lymphoma is highly associated with the non-fat portion of milk and inversely associated with whole grain cereals,²⁹ it is likely that calcium is involved in the etiology. It is not likely to be either lactose, which is implicated in ischemic heart disease,²⁸ or milk protein, which is implicated in osteoporosis,³⁰ both of which have lower associations with adult lymphoma than the non-fat portion of milk. Phytate in whole grains reduces calcium absorption,³¹ and whole grains also have other protective compounds.²⁹ Whole grain food intake has been found to be inversely related to lymphoma (OR = 0.5),³² while other researchers have found

Table 5: Ratio of Male Lymphoma Mortality Rates Divided by that for Females.

| Age range (years) | 55-64 | 65-74 | 75+ |
|--------------------|-------|-------|------|
| 1986 mean values | 1.52 | 1.55 | 1.58 |
| 1986 median values | 1.52 | 1.49 | 1.50 |
| 1994 mean values | 1.47 | 1.52 | 1.48 |
| 1994 median values | 1.51 | 1.45 | 1.43 |

refined-cereal intake to be positively associated with a number of cancers.³³ The ecological results presented in Table 4 did not find a statistically significant risk-reduction factor for whole and refined cereals combined. However, if whole grains alone could be studied using the ecological approach, a protective effect might well be found.

A number of studies have investigated the roles of vitamin D metabolism and hypercalcemia in relation to Hodgkin's lymphoma (HL). In one case, a patient with HL experienced abnormal vitamin D metabolism that resulted in hypercalcemia when he was exposed to ultraviolet radiation.³⁴ It was hypothesized that HL caused abnormal vitamin D metabolism. A number of such cases have been reviewed.³⁵ Calcitriol has been implicated as a hematolymphoid regulatory hormone.³⁶ It has been reported that deregulated synthesis of a $1,25(\text{OH})_2\text{-D}$ -like metabolite is a common cause of hypercalcemia and hypercalciuria in patients with lymphoma.³⁷ Abnormal calcium metabolism, hypercalciuria, and dysregulated calcitriol production are common findings in normocalcemic patients with NHL.³⁸ Adult T-cell leukemia-lymphoma patients with elevated serum calcium levels were found to have lower survival rates than those with normal serum calcium.³⁹ Researchers report that calcitriol-mediated A-to-B calcium transport in Caco-2 cells is a specific, transcellular process that requires transcriptional events normally mediated through the vitamin D receptor, with $1,25(\text{OH})_2\text{-D}_3$ being the regulator.⁴⁰ Cats with neoplasia, the most common of which were lymphoma and squamous cell carcinoma, had higher serum calcium levels than those with renal failure.⁴¹ Hepatic and serum levels of iron, zinc, and calcium were highly elevated 15 days after transplantation of Dalton's lymphoma in male Swiss albino mice.⁴² From monitoring gamma-glutamyl transferase levels in the liver as well, it was concluded that the metals studied may have a role in initiating and controlling cellular proliferation.

While these papers support the hypothesis that HL gives rise to vitamin D metabolism abnormalities followed by hypercalcemia, that may not be the complete story since excess dietary calcium seems to be linked to lymphoma mortality rates. It could be that the excess calcium from milk, with or without a disruption of vitamin D metabolism, is involved in the etiology of lymphoma among adults. The role of cereals may either be to displace milk from the diet or to reduce the absorption of calcium, perhaps through phytates found in whole grains, as discussed above.

The statistical results for the non-fat portion of milk given in Tables 2 and 3 can be used to estimate the fraction of lymphoma mortality due to diet. For the 1994 data, the mortality rate increased by a factor of 2.3 from age 55-64 years to age 65-74 years, and a factor of 1.8 from 65-74 years to 75+ years. Suitably averaging the r^2 values for the three age groups gives 0.69 for males and 0.71 for females. Thus, diet seems to explain approximately 70 percent of the lymphoma mortality rates on a global basis. On the other hand, the annual incidence of NHL has risen in the United States from 5.9 per 100,000 in 1950 to 13.7 in 1989, which may be consistent with chemical exposure increases during this period. Milk consumption was relatively unchanged during that period, with the per capita supply staying in the range 342-376 kCal/day from 1962 to 1993. However, cancers generally have several risk and risk-reduction factors, so the increase in incidence rates for NHL does not necessarily argue against the results of this study.

In another study,⁴³ the roles of solar UV-B exposure and vitamin D as risk reduction factors for various forms of cancer were studied. NHL and prostate cancer were found to have similar fractions of U.S. mortality from 1970-1994 related to insufficient solar UV-B radiation, 3-5 percent or 7-9 percent, respectively, depending on the UV-B data used, which was lower than for most other cancers

studied. This finding is consistent with a common etiology, dietary calcium in this case. Many of the other cancer types have dietary fat as the high risk factor.

Note that the dietary links to lymphoma differ from those for leukemia. Using the ecological approach, age-adjusted mortality data⁴⁴⁻⁴⁵ found total dietary fat to have the highest association with leukemia. However, using age-adjusted incidence data for those aged 35-64 years, it was found that total caloric supply gave the best results.⁴⁶ Using the 1986 leukemia mortality data¹⁹ it can be shown that leukemia mortality is most highly associated with total energy supply of adults under the age of 64 years, after which dietary fat has the highest association. This point is made here to demonstrate that leukemia and lymphoma have different dietary links, hence, different etiologies, and to suggest that the ecological approach is not just selecting out non-fat milk as a general risk for cancer mortality.

Whether disruptions in calcium metabolism are a cause or an effect of lymphoma (or both) still remains to be seen. It is hoped that the results presented here will lead to more case-control, clinical, and cohort studies to evaluate and extend these findings and result in guidelines for reducing the risk of adult lymphoma.

Acknowledgements

The author thanks the staffs of the Moorman Memorial Library of the Eastern Virginia Medical School (Norfolk, VA), the Health Sciences Library of the Riverside School of Professional Nursing (Newport News, VA), and the Tomkins-McCaw Library of the Virginia Commonwealth University (Richmond, VA) for the use of their facilities.

References

1. Hoar SK, Blair A, Holmes FF, et al. Agricultural herbicide use and risk of lymphoma and soft-tissue sarcoma. *JAMA* 1986;256:1141-1147.
2. Kogevinas M, Kauppinen T, Winkelmann R, et al. Soft tissue sarcoma and non-Hodgkin's lymphoma in workers exposed to phenoxy herbicides, chlorophenols, and dioxins: two nested case-control studies. *Epidemiology* 1995;6:396-402.
3. Cantor KP, Blair A, Everett G, et al. Pesticides and other agricultural risk factors for non-Hodgkin's lymphoma among men in Iowa and Minnesota. *Cancer Res* 1992;52:2447-2455.
4. Pearce N, Smith AH, Reif JS. Increased risks of soft tissue sarcoma, malignant lymphoma, and acute myeloid leukemia in abattoir workers. *Am J Ind Med* 1988;14:63-72.
5. La Vecchia C, Negri E, D'Avanzo B, Franceschi S. Occupation and lymphoid neoplasms. *Br J Cancer* 1989;60:385-388.
6. Franceschi S, Serraino D, La Vecchia C, et al. Occupation and risk of Hodgkin's disease in north-east Italy. *Int J Cancer* 1991;48:831-835.
7. Brown LM, Everett GD, Gibson R, et al. Smoking and risk of non-Hodgkin's lymphoma and multiple myeloma. *Cancer Causes Control* 1992;3:49-55.
8. De Stefani E, Fierro L, Barrios E, Ronco A. Tobacco, alcohol, diet and risk of non-Hodgkin's lymphoma: a case-control study in Uruguay. *Leuk Res* 1998;22:445-452.
9. Ursin G, Bjelke E, Heuch I, Vollste SE. Milk consumption and cancer incidence: a Norwegian prospective study. *Br J Cancer* 1990;61:456-459.
10. Franceschi S, Serraino D, Carbone A, et al. Dietary factors and non-Hodgkin's lymphoma: a case-control study in the northeastern part of Italy. *Nutr Cancer* 1989;12:333-341.
11. Ward MH, Zahm SH, Weisenburger DD, et al. Dietary factors and non-Hodgkin's lymphoma in Nebraska (United States). *Cancer Causes Control* 1994;5:422-432.
12. Tavani A, Pregnolato A, Negri E, et al. Diet and risk of lymphoid neoplasms and soft tissue sarcomas. *Nutr Cancer* 1997;27:256-260.
13. Chiu BC, Cerhan JR, Folsom AR, et al. Diet and risk of non-Hodgkin lymphoma in older women. *JAMA* 1996;275:1315-1321.
14. Zhang S, Hunter DJ, Rosner BA, et al. Dietary fat and protein in relation to risk of non-Hodgkin's lymphoma among women. *J Natl Cancer Inst* 1999;91:1751-1758.

15. Pryme IF, Bardocz S, Pusztai A, Ewen SW. The growth of an established murine non-Hodgkin lymphoma tumour is limited by switching to a phytohaemagglutinin-containing diet. *Cancer Lett* 1999;146:87-91.
16. Dockerty JD, Skegg DC, Elwood JM, et al. Infections, vaccinations, and the risk of childhood leukaemia. *Br J Cancer* 1999;80:1483-1489.
17. Severson RK, Ross JA. The causes of acute leukemia. *Curr Opin Oncol* 1999;11:20-24.
18. Shu XO, Linet MS, Steinbuch M, et al. Breast-feeding and risk of childhood acute leukemia. *J Natl Cancer Inst* 1999;91:1765-1772.
19. World Health Organization. *World Health Statistics Annual*, Geneva, 1986-1998.
20. Food and Agriculture Organization (FAO) of the United Nations, *Food Balance Sheets*, Rome, Italy, 1991.
21. Food and Agriculture Organization (FAO) of the United Nations, *Food Balance Sheets*, Rome, Italy, 1996.
22. U.S. Dept. of Agriculture, Agriculture Research Service. The 1994-96 Continuing Survey of Food Intakes by Individuals and the 1994-96 Diet and Health Knowledge Survey, CD-ROM, available from NTIS, Springfield, MD, 1998.
23. Sasaki S, Kesteloot H. Value of Food and Agriculture Organization data on food-balance sheets as a data source for dietary fat intake in epidemiologic studies. *Am J Clin Nutr* 1992;56:716-723.
24. Williams C, Wiseman M, Buttriss J. Food-based dietary guidelines—a staged approach. *Br J Nutr* 1999;81:S131-S153.
25. Becker W, Foley S, Shelley E, Gibney M. Energy under-reporting in Swedish and Irish dietary surveys: implications for food-based dietary guidelines. *Br J Nutr* 1999;81:S127-S131.
26. Jula A, Seppanen R, Alanen E. Influence of days of the week on reported food, macronutrient and alcohol intake among an adult population in south western Finland. *Eur J Clin Nutr* 1999;53:808-812.
27. World Health Organization, *Tobacco Alert*, Geneva, 1996.
28. Grant WB. Milk and other dietary influences on coronary heart disease. *Altern Med Rev* 1998;3:281-294.
29. Slavin J, Jacobs D, Marquart L. Whole-grain consumption and chronic disease: protective mechanisms. *Nutr Cancer* 1997;27:14-21.
30. Grant WB. Lactose maldigestion and calcium from dairy. *Am J Clin Nutr* 1999;70:301-302.
31. Weaver CM, Heaney RP, Martin BR, Fitzsimmons ML. Human calcium absorption from whole-wheat products. *J Nutr* 1991;21:1769-1775.
32. Chatenoud L, Tavani A, La Vecchia C, et al. Whole grain food intake and cancer risk. *Int J Cancer* 1998;77:24-28.
33. Chatenoud L, La Vecchia C, Franceschi S, et al. Refined-cereal intake and risk of selected cancers in Italy. *Am J Clin Nutr* 1999;70:1107-1110.
34. Davies M, Mawer EB, Hayes ME, Lumb GA. Abnormal vitamin D metabolism in Hodgkin's lymphoma. *Lancet* 1985;1:1186-1188.
35. Rieke JW, Donaldson SS, Horning SJ. Hypercalcemia and vitamin D metabolism in Hodgkin's disease. Is there an underlying immunoregulatory relationship? *Cancer* 1989;63:1700-1707.
36. Davies M, Hayes ME, Yin JA, et al. Abnormal synthesis of 1,25-dihydroxyvitamin D in patients with malignant lymphoma. *J Clin Endocrinol Metab* 1994;78:1202-1207.
37. Adams JS, Fernandez M, Gacad MA, et al. Vitamin D metabolite-mediated hypercalcemia and hypercalciuria patients with AIDS- and non-AIDS-associated lymphoma. *Blood* 1989;73:235-239.
38. Seymour JF, Gagel RF, Hagemeister FB, et al. Calcitriol production in hypercalcemic and normocalcemic patients with non-Hodgkin lymphoma. *Ann Intern Med* 1994;121:633-640.
39. Hoshi H, Nagamachi S, Jinnouchi S, et al. Bone scintigraphy as a prognostic factor in patients with adult T-cell leukemia-lymphoma. *Clin Nucl Med* 1994;19:992-995.
40. Fleet JC, Wood RJ. Specific 1,25(OH)₂D₃-mediated regulation of transcellular calcium transport in caco-2 cells. *Am J Physiol* 1999;276:G958-G964.
41. Savary KC, Price GS, Vaden SL. Hypercalcemia in cats: a retrospective study of 71 cases (1991-1997). *J Vet Intern Med* 2000;14:184-189.

42. Bishayee A, Kulkarni SP, Roy K, Chatterjee M. Alterations in total iron, zinc, and calcium levels and their influence on the hepatic activities of gamma-glutamyl transferase and glucose-6-phosphatase in the host bearing transplantable murine lymphoma. *Cancer Invest* 1998;16:231-236.
43. Grant WB, Slusser JR. An estimation of excess cancer mortality in the United States due to inadequate doses of solar ultraviolet-B radiation and vitamin D, in preparation.
44. Carroll KK, Khor HT. Dietary fat in relation to tumorigenesis. *Prog Biochem Pharmacol* 1975;10:308-353.
45. Carroll KK, Braden LM, Bell JA, Kalamegham R. Fat and cancer. *Cancer* 1986;58:1818-1825.
46. Hursting SD, Margolin BH, Switzer BR. Diet and human leukemia: an analysis of international data. *Prev Med* 1993;22:409-422.